

ON BRAIDED AND RIBBON UNITARY FUSION CATEGORIES

CÉSAR GALINDO

ABSTRACT. We prove that every braiding over a unitary fusion category is unitary and every unitary braided fusion category admits a unique unitary ribbon structure.

1. INTRODUCTION

The notion of ribbon (or premodular) unitary fusion categories is important for their applications to low-dimensional topology and topological quantum computation (see [8] and [9]). They also appear naturally as invariants in subfactor theory and quantum groups theory.

During the AIM conference on “Classifying Fusion Categories” (March 2012) a list of problem was posted, see <http://aimpl.org/fusioncat/>. In this note we answer the second question of Problem 3.3, posted by Zhenghan Wang. We prove that every braiding over a unitary fusion category is unitary and every unitary braided fusion category admits a unique unitary ribbon structure. As a consequence, if the underlying fusion category of a modular category is unitary, we may freely choose a braiding to obtain a unitary braided fusion category and then there is exactly one choice of a ribbon structure that will make the associated Rational Conformal Field Theory unitary.

Acknowledgements The author would like to thank Eric Rowell and Zhenghan Wang for useful comments.

2. PRELIMINARIES

In this note, we will use basic theory of fusion categories and braided fusion categories. For further details on these topics, we refer the reader to [3]. In this section we recall some definitions and results on unitary fusion categories. Much of the material which appears here can be found in [8].

2.1. Unitary fusion categories. A C^* -category \mathcal{D} is a \mathbb{C} -linear abelian category with an involutive antilinear contravariant endofunctor $*$ which is the identity on objects, the hom-spaces $\text{Hom}_{\mathcal{D}}(X, Y)$ are Hilbert spaces and the norms satisfy

$$\|fg\| \leq \|f\| \|g\|, \quad \|f^*f\| = \|f\|^2,$$

for all $f \in \text{Hom}_{\mathcal{D}}(X, Y), g \in \text{Hom}_{\mathcal{D}}(Y, Z)$, where f^* denote the image of f under $*$.

Let X and Y be objects in a C^* -category. A morphism $u : X \rightarrow Y$ is **unitary** if $uu^* = \text{id}_Y$ and $u^*u = \text{id}_X$. A morphism $a : X \rightarrow X$ is **self-adjoint** if $a^* = a$.

Date: September 11, 2012.

2010 Mathematics Subject Classification. 20F36, 16W30, 18D10.

Remark 2.1. Every isomorphism in a C^* -category has a polar decomposition, *i.e.*, if $f : X \rightarrow Y$ is an isomorphism, then $f = ua$ where $a : X \rightarrow X$ is self-adjoint and $u : X \rightarrow Y$ is unitary, see [1, Proposition 8].

A **unitary fusion category** is a fusion category \mathcal{C} , where \mathcal{C} is a C^* -category, the constraints are unitary and $(f \otimes g)^* = f^* \otimes g^*$, for every pair of morphisms f, g in \mathcal{C} .

Remark 2.2.

- (1) A unitary fusion category is a fusion category with addition structure. Hence, a fusion category could have more than one unitary structure. All examples known to the author admit a unique unitary structure. Moreover, in [4, Theorem 5.20] it was proved that every weakly group-theoretical fusion category admits a unique unitary structure.
- (2) If \mathcal{C} is a unitary fusion category, we can find basis such that the F -matrices $(F_l^{ijk})_{n,m} = F_{l;n,m}^{i,j,k}$ are unitary, where $\{F_{l;n,m}^{i,j,k}\}$ are the $6j$ -symbols (see [9] or [8] for the definition of $6j$ -symbols). Conversely, if for a fusion category \mathcal{C} it is possible to find basis such that the F -matrices $(F_l^{ijk})_{n,m} = F_{l;n,m}^{i,j,k}$ are unitary, then \mathcal{C} is a unitary fusion category. See [10, Section 4].

3. BRAIDING AND MODULAR STRUCTURES OVER UNITARY FUSION CATEGORIES ARE UNITARY

3.1. The center of a unitary fusion category. We shall recall the definition of the **center** $\mathcal{Z}(\mathcal{C})$ of a monoidal category \mathcal{C} , see [5, Chapter XIII]. The objects of $\mathcal{Z}(\mathcal{C})$ are pairs $(Y, c_{-,Y})$, where $Y \in \mathcal{C}$ and $c_{X,Y} : X \otimes Y \rightarrow Y \otimes X$ are isomorphisms natural in X satisfying $c_{X \otimes Y, Z} = (c_{X,Z} \otimes \text{id}_Y)(\text{id}_X \otimes c_{Y,Z})$ and $c_{I,Y} = \text{id}_Y$, for all $X, Y, Z \in \mathcal{C}$. A morphism $f : (X, c_{-,X}) \rightarrow (Y, c_{-,Y})$ is a morphism $f : X \rightarrow Y$ in \mathcal{C} such that $(f \otimes \text{id}_W)c_{W,X} = c_{W,Y}(\text{id}_W \otimes f)$ for all $W \in \mathcal{C}$.

The center is a braided monoidal category with structure given as follows:

- the tensor product is $(Y, c_{-,Y}) \otimes (Z, c_{-,Z}) = (Y \otimes Z, c_{-,Y \otimes Z})$, where
$$c_{X,Y \otimes Z} = (\text{id}_Y \otimes c_{X,Z})(c_{X,Y} \otimes \text{id}_Z) : X \otimes Y \otimes Z \rightarrow Y \otimes Z \otimes X,$$
for all $X \in \mathcal{C}$,
- the identity element is $(I, c_{-,I})$, $c_{Z,I} = \text{id}_Z$
- the braiding is given by the morphism $c_{X,Y}$.

If \mathcal{C} is a unitary fusion category the **unitary center** $\mathcal{Z}^*(\mathcal{C})$ is defined as the full tensor subcategory of $\mathcal{Z}(\mathcal{C})$, where $(X, c_{-,X}) \in \mathcal{Z}^*(\mathcal{C})$ if and only if $c_{W,X} : W \otimes X \rightarrow X \otimes W$ is unitary for all $W \in \mathcal{C}$.

Proposition 3.1. *Let \mathcal{C} be a unitary fusion category then $\mathcal{Z}^*(\mathcal{C}) = \mathcal{Z}(\mathcal{C})$.*

Proof. Let $(X, c_{-,X})$ be an object in $\mathcal{Z}(\mathcal{C})$. By [4, Proposition 5.24.] or [7, Theorem 6.4], the inclusion functor $\mathcal{Z}^*(\mathcal{C}) \subseteq \mathcal{Z}(\mathcal{C})$ is a tensor equivalence. Therefore, there is an object $(Y, c_{-,Y})$ in $\mathcal{Z}^*(\mathcal{C})$ and an isomorphism $f : (X, c_{-,X}) \rightarrow (Y, c_{-,Y})$ in $\mathcal{Z}(\mathcal{C})$. By Remark 2.1 there exists a unitary arrow $u : (X, c_{-,X}) \rightarrow (Y, c_{-,Y})$. Hence, for every $W \in \mathcal{C}$, $c_{W,X} = (u \otimes \text{id}_W)^* \circ c_{W,Y} \circ (\text{id}_W \otimes u)$, so $c_{W,X}$ is an unitary arrow and $(X, c_{-,X}) \in \mathcal{Z}^*(\mathcal{C})$. \square

A braiding over a unitary fusion category \mathcal{C} is called **unitary braiding** if the morphism $c_{X,Y}$ is unitary for any pair of objects $X, Y \in \mathcal{C}$.

Theorem 3.2. *Every braiding of a unitary fusion category is unitary.*

Proof. Let \mathcal{C} be a unitary fusion category and let c be a braiding. It is easy to see that the braiding c defines an inclusion functor $\mathcal{C} \hookrightarrow \mathcal{Z}(\mathcal{C})$, $X \mapsto (X, c_X, -)$. Proposition 3.1 implies that $c_{X,W}$ is unitary for every $W \in \mathcal{C}$. \square

Remark 3.3.

- (1) Theorem 3.2 implies that if the F -matrices $(F_l^{ijk})_{n,m} = F_{l;n,m}^{i,j,k}$ are unitary, then the R -matrices of the braiding are always unitarily diagonalizable.
- (2) A Kac algebra $(H, m, \Delta, *)$ is a semisimple Hopf algebra such that $(H, *)$ is a C^* -algebra and the maps Δ and ε are C^* -algebra maps. Theorem 3.2 implies that every R -matrix in a Kac algebra is unitary in the sense that $R^* = R^{-1}$.

3.2. Ribbon structures on unitary fusion categories. If \mathcal{C} is a fusion category, then for every $f \in \text{Hom}_{\mathcal{C}}(X, Y)$ the **transpose** of f , is defined by ${}^t f := (X^* \otimes \text{ev}_Y)(X^* \otimes f \otimes Y^*)(\text{coev}_X \otimes Y^*) \in \text{Hom}_{\mathcal{C}}(Y^*, X^*)$.

A **twist** on a braided fusion category \mathcal{C} is a natural automorphism of the identity functor $\theta \in \text{Aut}(\text{Id}_{\mathcal{C}})$, such that

$$\theta_{X \otimes Y} = (\theta_X \otimes \theta_Y) c_{Y,X} c_{X,Y}$$

for all $X, Y \in \mathcal{C}$. A twist is called a **ribbon structure** if ${}^t \theta_X = \theta_{X^*}$. A fusion category with a ribbon structure is called a **ribbon fusion category**. Each ribbon structure θ defines a **quantum dimension function** by $\dim_{\theta}(X) = \text{ev}_X c_{X,X^*}(\theta_X \otimes \text{id}_{X^*}) \text{coev}_X$.

We shall denote by $\text{Aut}_{\otimes}(\text{Id}_{\mathcal{C}})_{(+,-)}$ the abelian group of tensor automorphisms of the identity γ such that $\gamma_X = \pm \text{id}_X$ for every simple object $X \in \mathcal{C}$.

Proposition 3.4. *Let \mathcal{C} be a braided fusion category. If the set of ribbon structures is not empty, it is a torsor under $\text{Aut}_{\otimes}(\text{Id}_{\mathcal{C}})_{(+,-)}$.*

Proof. Let θ and θ' be ribbon structures. It is easy to see that $\gamma := \theta^{-1} \circ \theta' : \text{Id}_{\mathcal{C}} \rightarrow \text{Id}_{\mathcal{C}}$ is a tensor automorphism of the identity. We may and shall assume that \mathcal{C} is skeletal. Then, $X = X^{**}$ and for every simple object, we have $\theta_X = \theta(X) \text{id}_X$, $\theta'_X = \theta'(X) \text{id}_X$, $\gamma_X = \gamma(X) \text{id}_X$ for some $\gamma(X), \theta(X), \theta'(X) \in \mathbb{C}^*$ and $\theta(X)' = \gamma(X) \theta(X)$. Since θ' is a ribbon structure, for every simple object $X \in \mathcal{C}$, $\dim_{\theta'}(X) = \dim_{\theta'}(X^*)$. On the other hand, $\dim_{\theta'}(X) = \gamma(X) \dim_{\theta}(X)$. Therefore $\gamma(X) = \gamma(X^*)$ and, since $\gamma(X^*) = \gamma(X)^{-1}$ we conclude that γ has order two.

Conversely, if γ is an automorphism of the identity such that $\gamma_X = \pm \text{id}_X$ for every simple object, then, for every ribbon structure θ , the natural isomorphism $\theta' = \theta \gamma$ is a new ribbon structure. \square

If \mathcal{C} is a unitary fusion category a ribbon structure on \mathcal{C} is called **unitary ribbon structure** if θ_X is unitary, $(\text{coev}_X)^* = \text{ev}_X \circ c_{X,X^*} \circ (\theta_X \otimes \text{id}_{X^*})$ and $(\text{ev}_X)^* = (\text{id}_{X^*} \otimes \theta_X^{-1}) \circ c_{X^*,X}^{-1} \circ \text{coev}_X$ for all $X \in \mathcal{C}$. A unitary fusion category with a unitary ribbon structure is called a **unitary ribbon fusion category**. In a unitary ribbon fusion category

$$\dim_{\theta}(X) = \text{ev}_X \circ c_{X,X^*} \circ (\theta_X \otimes \text{id}_{X^*}) \circ \text{coev}_X = (\text{coev}_X)^* \circ \text{coev}_X,$$

therefore, the quantum dimension of every object is a positive number.

Theorem 3.5. *Every braided fusion category with a unitary structure admits a unique unitary ribbon structure.*

Proof. By [6, Proposition 2.4] every braided unitary fusion category admits a canonical unitary ribbon structure. Let θ_c the canonical ribbon structure associated to c . By Proposition 3.4, if θ' is another unitary ribbon structure, then there is $\gamma \in \text{Aut}_{\otimes}(\text{Id}_{\mathcal{C}})_{(+,-)}$ such that $\theta' = \theta_c \gamma$. If γ is not the identity there is a simple object $X \in \mathcal{C}$ such that $\gamma_X = -\text{id}_X$, then $\dim_{\theta'}(X) = -\dim_{\theta_c}(X) < 0$, but the quantum dimension of every object of any unitary ribbon structure is positive. Therefore γ is the identity and θ_c is unique. \square

Remark 3.6. It follows from Theorem 3.5 that if a unitary braided fusion category is non-degenerate (see [2] for a definition), then it admits a unique unitary modular structure.

REFERENCES

- [1] J. Baez, “Higher-Dimensional Algebra II. 2-Hilbert Spaces.” *Adv. Math.* **127** (1997), 125–189.
- [2] V. Drinfeld, S. Gelaki, D. Nikshych and V. Ostrik, “On Braided Fusion Categories I.” *Selecta Math. N.S.* **16**, 1 (2010) 1–119.
- [3] P. Etingof, D. Nikshych and V. Ostrik, “On fusion categories.” *Ann. of Math.* (2) **162** (2005), no. 2, 581–642.
- [4] C. Galindo, S.-M. Hong, E. Rowell, “Generalized and quasi-localizations of braid group representations”, to appear in *Int. Math. Res. Not.*, arXiv:1105.5048.
- [5] C. Kassel, *Quantum Groups*, Graduate Texts in Mathematics **155**, Springer-Verlag, New York, 1995.
- [6] M. Mueger, “Galois theory for braided tensor categories and the modular closure.” *Adv. Math.* **150** (2000), 151–201.
- [7] M. Mueger, “From subfactors to categories and topology II. The quantum double of tensor categories and subfactors.” *J. Pure Appl. Alg.* **180** (2003), 159–219.
- [8] V. G. Turaev, *Quantum Invariants of Knots and 3-Manifolds.*, Revised Edition, de Gruyter Studies in Mathematics **18**, Walter de Gruyter & Co., Berlin, 2010.
- [9] Z. Wang, *Topological quantum computation.*, CBMS Regional Conference Series in Mathematics, **112**, 2010.
- [10] S. Yamagami, “Polygonal presentations of semisimple tensor categories.”, *J. Math. Soc. Japan*, **54**, (2002), 61–88.

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DE LOS ANDES,
BOGOTÁ, COLOMBIA
E-mail address: cn.galindo1116@uniandes.edu.co, cesarneyit@gmail.com